

Claims:

Sub 1
5 1. A pharmaceutical preparation which provides a dosage form of Tamoxifen, ~~characterised in that~~ ^{wherein} the dosage form is that of a solution in a suitable solvent of at least 1.5 g/ml of Tamoxifen Citrate.

2. A pharmaceutical preparation according to claim 1, characterised in that the solvent comprises the following components: (a) of from 10% to 20% by weight of ethanol; (b) of from 10% to 60% by weight of a glycol; and (c) water, optionally containing additives, in a volume percentage adding up to 100% by volume.

3. A pharmaceutical preparation according to claim 2, ~~characterised in that~~ ^{wherein} the glycol component is a mixture of propylene glycol and glycerol.

15 4. A pharmaceutical preparation according to claim 3, ~~characterised in that~~ ^{wherein} the water component (c) contains a bulk-sweetening agent.

5. A pharmaceutical preparation according to claim 4, ~~characterised in that~~ ^{wherein} the bulk-sweetening agent is ~~of~~ from 15% to 25 % by weight of sorbitol.

20 6. A pharmaceutical preparation according to claim 5, ~~characterised in that~~ ^{wherein} the solvent comprises the following components: 15% by weight of ethanol, 10% by weight of propylene glycol, 50% by weight of glycerol, 20% by weight of a solution of 70% by weight of sorbitol in water, and water, in a volume percentage adding up to 100% by volume.

25 7. A process for the preparation of a solution according to claim 2, comprising the steps of dissolving the Tamoxifen Citrate in ~~the mixed~~ ^{a mixture of the} ethanol and glycol components and then adding the other components.

9

8. A process for the preparation of a solution according to claim 2, comprising the steps of first dispersing Tamoxifen Citrate in the glycol component and then adding the ethanol component and the water component.

5

9. A process according to claim 7, characterised in that a solution in accordance with claim 3 is made by first dispersing Tamoxifen Citrate in the propylene glycol to form a dispersion, adding dispersion to the glycerol and, then adding the ethanol component to form a solution.

Sub As

10

Add As

10